



Acetaminophen (Tylenol®) Dosage Chart

Choose the dose closest to your child's weight and give every 4-6 hours as needed.

Weight		Dose 15 mg/kg	Oral Suspension 160 mg/5ml	
pounds	kilograms	mg	ml	teaspoons
6-11	2.7 - 5.3	40 mg	1.25 ml	$\frac{1}{4}$ tsp
12 - 17	5.4 - 8.1	80 mg	2.5 ml	$\frac{1}{2}$ tsp
18 - 23	8.2 - 10.8	120 mg	3.75 ml	$\frac{3}{4}$ tsp
24 - 35	10.9 - 16.3	160 mg	5 ml	1 tsp
36 - 47	16.4 - 21.7	240 mg	7.5 ml	1 $\frac{1}{2}$ tsp
48 - 59	21.8 - 27.2	320 mg	10 ml	2 tsp
60 - 71	27.3 - 32.6	400 mg	12.5 ml	2 $\frac{1}{2}$ tsp
72 - 95	32.7 - 43.2	480 mg	15 ml	3 tsp

Ibuprofen (Motrin® or Advil®) Dosage Chart

Important: Do not give to infants under 6 months of age.

Choose the dose closest to your child's weight and give every 6-8 hours as needed.

Weight		Dose 10 mg/kg	Concentrated Drops 50 mg/1.25 ml	Oral Suspension 100 mg/5 ml	
pounds	kilograms	mg	ml	ml	teaspoons
12 - 17	5.4 - 8.1	50 mg	1.25 ml	2.5 ml	$\frac{1}{2}$ tsp
18 - 23	8.2 - 10.8	75 mg	1.875 ml	3.75 ml	$\frac{3}{4}$ tsp
24 - 35	10.9 - 16.3	100 mg	2.5 ml	5 ml	1 tsp
36 - 47	16.4 - 21.7	150 mg		7.5 ml	1 $\frac{1}{2}$ tsp
48 - 59	21.8 - 27.2	200 mg		10 ml	2 tsp
60 - 71	27.3 - 32.6	250 mg		12.5 ml	2 $\frac{1}{2}$ tsp
72 - 95	32.7 - 43.2	300 mg		15 ml	3 tsp

Diphenhydramine (Benadryl®) Dosage Chart

Choose the dose closest to your child's weight and give every 6 hours as needed.

Weight	Children's Allergy Liquid 12.5 mg = 5 ml		Chewable Tablets 12.5 mg each	Tablets 25 mg each
pounds	ml	teaspoons	Chewable tablets	tablets
11 - 16	2.5 ml	$\frac{1}{2}$	Use liquid	Use liquid
17 - 21	3.75 ml	$\frac{3}{4}$	Use liquid	Use liquid
22 - 32	5 ml	1	1	Use liquid or chewables
33 - 42	7.5 ml	1 $\frac{1}{2}$	1 $\frac{1}{2}$	Use liquid or chewables
43 - 53	10 ml	2	2	1
54 - 64	12.5 ml	2 $\frac{1}{2}$	2 $\frac{1}{2}$	1
65 - 75	15 ml	3	3	1 $\frac{1}{2}$
76 - 86	17.5 ml	3 $\frac{1}{2}$	3 $\frac{1}{2}$	1 $\frac{1}{2}$
> 86	20 ml	4	4	2