

## IMMUNIZATION AND WELL VISIT SCHEDULE

| Well Check Visit                  | What happens at this visit?   |
|-----------------------------------|---|
| 1-3 days after hospital discharge | 1st office visit--Newborn Well Check  |
| 2 weeks from date of birth        | Repeat Newborn Metabolic Screen (1st blood test done at hospital),<br>Edinburgh postpartum screen questionnaire           |
| 2 months                          | VAXELIS (DTaP/IPV/Hib/Hep B), Rotateq (oral),<br>VAXNEUVANCE (PCV15), Edinburgh postpartum screen<br>questionnaire        |
| 4 months                          | VAXELIS (DTaP/IPV/Hib/Hep B), Rotateq (oral),<br>VAXNEUVANCE (PCV15), TB screen questionnaire                             |
| 6 months                          | VAXELIS (DTaP/IPV/Hib/Hep B), Rotateq (oral),<br>VAXNEUVANCE (PCV15), <b>INFLUENZA (seasonal)</b> ,<br>Lead questionnaire |
| 9 months                          | Developmental Screening (PEDS:DM™)  |
| 12 months                         | MMR, Varivax, Hep A, Vision Spot Screen,<br>Anemia Check (blood test)   |
| 15 months                         | Pentacel (DTaP/IPV/Hib), VAXNEUVANCE (PCV15)  |
| 18 months                         | Hep A, Proquad (MMR/Varicella),<br>Autism Screening (M-CHAT-R) &<br>Developmental Screening (PEDS:DM™)                    |
| 24 months                         | Developmental Screening (M-CHAT-R), Vision Spot Screen  |
| 30 months                         | Developmental Screening (PEDS:DM™)  |
| 3 years                           | Vision Spot Screening   |
| 4 years                           | Quadracel (DTaP/IPV),<br>Proquad (MMR/Varicella) if not done at 18mos.,<br>Vision & Hearing                               |
| 3-18 years                        |   |
| 9 years                           | Well Child Check every year<br>(vision & hearing every other year)  |
| 10 years                          | Cholesterol Screen, HPV#1 (2 dose schedule start before 15yrs   |
| 11 years                          | old) HPV #2, Vision & Hearing screen  |
| 16 years                          | Tdap, MenQuadfi (MCV4)  |
| 17-18 years                       | MenQuadfi (MCV4) dose #2, Meningitis B dose #1 (Bexsero)  |
|                                   | Meningitis B dose #2, cholesterol screen  |